



Multiple solutions for a second order equation with radiation boundary conditions

Pablo Amster^{1, 2} and Mariel Paula Kuna ^{1, 2}

¹Departamento de Matemática, Facultad de Ciencias Exactas y Naturales,
Universidad de Buenos Aires, Argentina

²IMAS - CONICET, Ciudad Universitaria, Pabellón I, (1428) Buenos Aires, Argentina

Received 5 September 2016, appeared 11 May 2017

Communicated by Gennaro Infante

Abstract. A second order ordinary differential equation with a superlinear term is studied under radiation boundary conditions. Employing the variational method and an accurate shooting-type argument, we prove the existence of at least three or five solutions, depending on the interaction of the nonlinearity with the spectrum of the associated linear operator and the values of the radiation parameters.

Keywords: second order ODEs, radiation boundary conditions, multiplicity of solutions, variational method, shooting method.

2010 Mathematics Subject Classification: 34B15, 35A15.

1 Introduction

In a recent paper [1], the following problem was studied

$$u''(x) = g(x, u(x)) + A(x) \quad (1.1)$$

under radiation boundary conditions

$$u'(0) = a_0 u(0), \quad u'(1) = a_1 u(1). \quad (1.2)$$

Unlike the standard Robin condition, both coefficients a_0 and a_1 in the radiation boundary condition (1.2) are assumed to be strictly positive. Here, $A \in C([0, 1])$ and $g : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ is continuous, of class C^1 with respect to u and superlinear, that is:

$$\lim_{|u| \rightarrow +\infty} \frac{g(x, u)}{u} = +\infty \quad (1.3)$$

uniformly in $x \in [0, 1]$. Without loss of generality, it is assumed that $g(x, 0) = 0$ for all $x \in [0, 1]$.

 Corresponding author. Emails: pamster@dm.uba.ar (P. Amster), mpkuna@dm.uba.ar (M. P. Kuna)

The previous problem was motivated as a generalization of the following Painlevé II model for two-ion electrodiffusion studied in [2],

$$u''(x) = Ku(x)^3 + L(x)u(x) + A, \quad (1.4)$$

with K and A positive constants and $L(x) := a_0^2 + (a_1^2 - a_0^2)x$. As readily observed, the associated functional \mathcal{J} is coercive over a subspace $H \subset H^1(0,1)$ of codimension 1 and $-\mathcal{J}$ is coercive over a linear complement of H . This geometry explains the nature of the results in [2], where it was shown that the functional is in fact coercive over the whole space and, consequently, it achieves a global minimum but, under appropriate conditions, it admits also a local minimum and a saddle type critical point. In more precise terms, it was proved that the global minimizer of \mathcal{J} corresponds to a negative solution of the problem; moreover, if $a_1 \leq a_0$ then there are no other solutions. When $a_1 > a_0$, the solution is still unique for $A \gg 0$ but, when A is sufficiently small, the problem has at least three solutions.

With these ideas in mind, an extension of the above mentioned results was obtained in [1] for a general superlinear g nondecreasing in u and $A \geq 0$; namely, a solution always exists and, moreover, it is unique if A is sufficiently large or if $\frac{\partial g}{\partial u}(\cdot, u) > -\lambda_1$ for all u , where λ_1 is the first eigenvalue of the linear operator $Lu := -u''$ under the boundary condition (1.2). When A is close to 0, the problem has at least three solutions if $\frac{\partial g}{\partial u}(\cdot, 0) \leq -\lambda_1$. Furthermore, under an extra assumption, which is fulfilled for the particular case (1.4), the previous multiplicity result is sharp, in the sense that the number of solutions is *exactly* equal to 3. As a corollary, the mentioned results yield, for arbitrary A , the existence of at least three solutions when a_1 is large and a unique solution when a_1 is small.

The present paper is devoted to obtain further generalizations of the results in [1], by dropping the monotonicity assumption for g . After deducing some basic properties on the spectrum and eigenfunctions of the operator L , in the two first results we shall assume that $-\frac{\partial g}{\partial u}(x, 0)$ lies between consecutive eigenvalues, that is:

$$-\lambda_k \geq \frac{\partial g}{\partial u}(\cdot, 0) \geq -\lambda_{k+1} \quad (1.5)$$

where $\lambda_1 < \lambda_2 < \dots \rightarrow +\infty$ are the eigenvalues of L and, for convenience, we denote $\lambda_0 := -\infty$. As we shall see, λ_2 is always positive, so Theorem 2.9 in [1] is a direct consequence of the following theorem with $k = 1$.

Theorem 1.1. *Let (1.3) hold and assume that (1.5) is satisfied for some odd k . Then there exists $A_1 > 0$ such that (1.1)–(1.2) admits at least three solutions for $\|A\|_\infty < A_1$.*

In order to study the case in which k is even, it is worth recalling, from the mentioned uniqueness result in [1], that multiple solutions cannot be expected under the sole assumption that (1.5) holds for $k = 0$. In this sense, the following result can be regarded as complementary to Theorem 1.1 and allows to gain more solutions provided that $\lambda_1 < 0$. As we shall see, the latter condition is equivalent to a largeness condition on a_1 .

Theorem 1.2. *Let (1.3) hold and assume that $a_1 > \frac{a_0}{a_0+1}$. If (1.5) is satisfied for some even $k > 0$, then there exists $A_1 > 0$ such that (1.1)–(1.2) has at least five solutions for $\|A\|_\infty < A_1$.*

The preceding multiplicity results shall be proved by a shooting-type argument. It is worthy noticing, however, that solutions of the initial value problem typically blow up before $x = 1$ for large values of the shooting parameter λ . In order to overcome this difficulty, we

shall define an endpoint function that allows to reduce the problem to an equivalent one, with λ belonging to some appropriate interval $[-M, M]$. Once that a shooting operator T is established, the existence of multiple solutions is deduced by studying the sign changes of T . With this aim, we shall prove a fundamental lemma concerning the linearised problem at $u = 0$, which yields a straightforward proof of Theorem 1.1. Indeed, for an appropriate value $M > 0$ and $A = 0$, it shall be seen that $T(M) > 0 > T(-M)$ and $T(0) = 0$. Moreover, assumption (1.5) with k odd implies that $T'(0) < 0$; thus, by a continuity argument it shall be deduced that T has at least three roots when A is small. The situation in Theorem 1.2 is different because if k is even then $T'(0) > 0$; however, employing the extra assumption $\lambda_1 < 0$ we shall adapt the method of upper and lower solutions in order to obtain values $\lambda_- < 0 < \lambda_+$ such that $T(\lambda_-) > 0 > T(\lambda_+)$. This ensures that T has at least five roots when A is small. It is interesting to observe that some of the conclusions can be extended to more general situations, e.g. a system of equations, by the use of topological degree, although the results for the scalar case are sharper and, for this reason, deserve to be treated separately. Generalizations for systems of equations and other situations shall take part of a forthcoming paper.

Variational methods allow to obtain multiple solutions under a different condition, which is weaker than (1.5), by imposing a lower bound on the primitive of g , namely $G(x, u) := \int_0^u g(t, x) ds$. In order to formulate a precise statement, let us denote by φ_k the (unique) eigenfunction associated to λ_k such that $\varphi_k(0) > 0$ and $\|\varphi_k\|_{L^2} = 1$ and observe that, by superlinearity, the function G achieves a (nonpositive) minimum.

Theorem 1.3. *Assume there exist $K > 0$ and $k \in \mathbb{N}$ such that*

$$\frac{\partial g}{\partial u}(x, u) < -\lambda_k + 2 \frac{\|\varphi_k\|_{\infty}^2}{K^2} \min G \quad (1.6)$$

for all x and $|u| \leq K$. Then there exists a constant $A_1 > 0$ such that problem (1.1)–(1.2) has at least three solutions for $\|A\|_{L^2} < A_1$.

Our last result is devoted to analyse uniqueness and multiplicity according to the different values of the parameter a_1 . As we shall see, if a_1 is large then $\lambda_1 \ll 0$; thus, condition (1.6) is fulfilled with $k = 1$. Further considerations will show that the value A_1 in Theorem 1.3 can be made arbitrarily large as a_1 increases, yielding the existence of at least three solutions. The situation is different when $a_1 \rightarrow 0^+$, because λ_1 tends to some positive constant and the validity of conditions like (1.5) or (1.6) depends on the choice of g . Indeed, for a_1 arbitrarily small it is possible to find g and A such that the problem admits three solutions; however, if the derivative of g lies always above an appropriate constant, then the uniqueness condition given in [1] holds for a_1 small. More precisely, the following theorem holds.

Theorem 1.4. *There exists a constant a^* (depending only on $\|A\|_{L^2}$) such that problem (1.1)–(1.2) has at least three solutions for $a_1 > a^*$. Assume, moreover, that $\frac{\partial g}{\partial u}(\cdot, u) \geq c > -r_1^2$ for all u where $r_1 \in (0, \frac{\pi}{2})$ is the unique value such that $r_1 \tan r_1 = a_0$. Then there exists $a_* > 0$ such that the solution of (1.1)–(1.2) is unique when $0 < a_1 < a_*$.*

The paper is organized as follows. In the next section, we shall prove the basic facts concerning the eigenvalues of the linear operator L under the radiation boundary conditions that shall be used in the proofs of our main results. In Section 3, we define an accurate shooting-type operator and prove two existence results from which Theorems 1.1 and 1.2 are deduced in a straightforward manner. Finally, in Section 4 we shall apply a variational method in order to prove a quantitative version of Theorem 1.3, which provides a lower bound for A_1 and yields a proof of Theorem 1.4.

2 Spectrum of the associated linear operator

In this section, we shall obtain some elementary properties of the spectrum of the operator $Lu := -u''$, which is symmetric under the boundary condition (1.2). It is readily seen that all the eigenvalues of L are simple and form a sequence $\lambda_1 < \lambda_2 < \dots \rightarrow +\infty$. Zeros of an arbitrary eigenfunction φ are obviously simple; in particular, from the boundary condition it is deduced that φ does not vanish on the boundary. As mentioned, we shall denote $\lambda_0 := -\infty$ and, for $k > 0$, we shall set φ_k as the unique eigenfunction associated to λ_k such that $\varphi_k(0) > 0$ and $\|\varphi_k\|_{L^2} = 1$. Thus, $\{\varphi_j\}_{j \geq 1}$ is an orthonormal basis of $L^2(0, 1)$. A standard argument shows that φ_1 does not vanish and, by comparison, the k -th eigenfunction φ_k has exactly $k - 1$ zeros in $(0, 1)$. This, in turn, implies $\text{sgn}(\varphi_k(1)) = (-1)^{k-1}$.

Lemma 2.1. *The following properties hold:*

1. λ_1 is a (continuous) strictly decreasing function of a_1 ;
2. $\lambda_1 = 0$ if and only if $a_1 = \frac{a_0}{a_0+1}$;
3. $\lambda_1 < -a_1^2$ if and only if $a_1 > a_0$;
4. $\lambda_2 > 0$.

Proof. Let $a_1 < \tilde{a}_1$ and consider the respective eigenvalues and eigenfunctions $\lambda_1, \tilde{\lambda}_1$ and $\varphi_1, \tilde{\varphi}_1 > 0$. If $\lambda_1 \leq \tilde{\lambda}_1$, then

$$-\varphi_1'' \tilde{\varphi}_1 = \lambda_1 \varphi_1 \tilde{\varphi}_1 \leq \tilde{\lambda}_1 \varphi_1 \tilde{\varphi}_1 = -\tilde{\varphi}_1'' \varphi_1$$

and, after integration,

$$-a_1 \varphi_1(1) \tilde{\varphi}_1(1) \leq -\tilde{a}_1 \tilde{\varphi}_1(1) \varphi_1(1),$$

a contradiction. Continuity of λ_1 is left as an exercise for the reader. Moreover, a simple computation shows that 0 is eigenvalue if and only if $a_1 = \frac{a_0}{a_0+1}$; in this case, the corresponding eigenfunction is a linear function which, due to the boundary condition, cannot change sign and hence $0 = \lambda_1$. As a consequence, we deduce that $\lambda_1 < 0$ if and only if $a_1 > \frac{a_0}{a_0+1}$, in which case we may write $\lambda_1 = -r^2 < 0$, where $r > 0$ satisfies

$$\frac{r - a_0}{r + a_0} = \frac{r - a_1}{r + a_1} e^{2r}.$$

In particular, this shows that $\lambda_1 < -a_1^2$ if and only if $a_1 > a_0$. Finally, let us show that the second eigenvalue is always positive: indeed, otherwise $\lambda_2 < 0$, which implies $\text{sgn}(\varphi_2''(x)) = \text{sgn}(\varphi_2(x))$ for all x such that $\varphi_2(x) \neq 0$. From the boundary condition, we deduce that φ_2 does not vanish in $[0, 1]$, a contradiction. \square

The next lemma shall be the key for our proofs of Theorems 1.1 and 1.2.

Lemma 2.2. *Let $a \in C([0, 1])$ satisfy $\lambda_k \leq a \leq \lambda_{k+1}$. If u is the unique solution of the initial value problem*

$$u''(x) + a(x)u(x) = 0, \quad u'(0) = a_0 u(0) = a_0, \quad (2.1)$$

then $\text{sgn}[u'(1) - a_1 u(1)] = (-1)^k$.

Proof. Let us firstly prove that $u'(1) \neq a_1 u(1)$. With this aim, set $X \subset H^2(0,1)$ as the set of those functions satisfying (1.2) and define the symmetric bilinear form given by

$$B(u, v) := - \int_0^1 (u''(x) + a(x)u(x))v(x) dx.$$

Let $X_k := \text{span}\{\varphi_1, \dots, \varphi_k\}$. If $u \in X_k \setminus \{0\}$, then we may write $u = \sum_{j=1}^k s_j \varphi_j$ and $-u'' = \sum_{j=1}^k s_j \lambda_j \varphi_j$. Thus,

$$B(u, u) = \sum_{j=1}^k s_j^2 \lambda_j - \int_0^1 a(x)u(x)^2 dx \geq \int_0^1 [\lambda_k - a(x)]u(x)^2 dx \geq 0.$$

Moreover, if $B(u, u) = 0$ then $s_j = 0$ for all $j < k$ and

$$\int_0^1 [\lambda_k - a(x)]\varphi_k(x)^2 dx = 0$$

and hence φ_k vanishes over a non-zero measure interval I , a contradiction. In the same way, if $Y_k \subset H$ is the subspace spanned by $\{\varphi_j\}_{j>k}$, then $B(u, u) < 0$ for all $u \in Y_k \setminus \{0\}$. If $u(1) = a_1 u(1)$, then $u \in X$ and $B(u, v) = 0$ for all v . From [4, Lemma 1] we deduce that $u = 0$, a contradiction.

Next, define $A_k := \{a \in C([0,1]) : \lambda_k \leq a \leq \lambda_{k+1}\}$ and $T : A_k \rightarrow \mathbb{R}$ given by $T(a) := u'(1) - a_1 u(1)$, where $u = u_a$ is defined by (2.1). Observe that T is continuous and does not vanish. Moreover, A_k is connected (for example, because it is convex); thus the sign of T is constant over A_k and we may assume that a is constant. Hence we may take, for each k , the first (in fact, unique) value $a \in (\lambda_k, \lambda_{k+1})$ such that $u_a(1) = 0$. It follows that $u'_a(1)$ and $\varphi_k(1)$ have opposite signs; thus, $\text{sgn}(T(a)) = \text{sgn}(u'_a(1)) = (-1)^k$. \square

3 Shooting method revisited

Let us recall that the multiplicity results in [1] were obtained from the application of a shooting-type method. However, the success of this procedure was strongly based on the monotonicity of g , which was employed to guarantee that the graphs of two different solutions of (1.1) satisfying the first condition in (1.2) do not intersect. This property does not hold for the general case, so it is required to define the shooting operator in a more careful way.

With this aim observe, in the first place, that solutions of (1.1)–(1.2) are bounded. This fact was proved in [1] although, for the sake of completeness, a short proof is sketched here. Let u be a solution and $\psi(x) := (a_1 - a_0)x + a_0$. Multiply by u and integrate to obtain, for some constant C :

$$\begin{aligned} \int_0^1 [u'(x)^2 + g(x, u(x))u(x) + A(x)u(x)] dx &= a_1 u(1)^2 - a_0 u(0)^2 \\ &= \int_0^1 [\psi'(x)u(x)^2 + \psi(x)2u(x)u'(x)] dx \leq C \|u\|_{L^2}^2 + \frac{1}{2} \|u'\|_{L^2}^2. \end{aligned}$$

Next, choose a constant K such that $\int_0^1 [g(\cdot, u)u + Au] \geq (C + \frac{1}{2}) \|u\|_{L^2}^2 - K$, then $\|u\|_\infty \leq \|u\|_{H^1} \leq \sqrt{2K}$ and so completes the proof.

In order to define our shooting operator, let us fix a constant $M > \sqrt{2K}$ such that

$$\frac{g(x, u) + A(x)}{u} > R$$

for $|u| \geq M$ and some R to be specified. For each $\lambda \in \mathbb{R}$, let u_λ be the unique solution of (1.1) with initial condition $u'(0) = a_0 u(0) = a_0 \lambda$. If $|\lambda| \leq M$ and $|u_\lambda|$ reaches the value $2M$ for the first time at some t_1 , then we may fix $t_0 < t_1$ such that $|u_\lambda(t_0)| = M$ and $M < |u_\lambda| < 2M$ over (t_0, t_1) . Since $\frac{u''_\lambda}{u_\lambda} > R$ and $u'_\lambda(t_0)u_\lambda(t_0) \geq 0$, it is deduced that $u''_\lambda u'_\lambda > R u_\lambda u'_\lambda$ over (t_0, t_1) , whence

$$u'_\lambda(t_1)^2 > u'_\lambda(t_0)^2 + 3RM^2 > a_1^2 u_\lambda(t_1)^2$$

provided that $R > \frac{4}{3}a_1^2$. This implies, on the one hand, that the 'endpoint' function

$$e(\lambda) := \begin{cases} t_1 & \text{if } t_1 \text{ exists} \\ 1 & \text{otherwise} \end{cases}$$

is continuous. On the other hand, the (continuous) function $T : [-M, M] \rightarrow \mathbb{R}$ given by

$$T(\lambda) := u'_\lambda(e(\lambda)) - a_1 u_\lambda(e(\lambda))$$

characterizes the solutions of (1.1)-(1.2), in the sense that u is a solution if and only if there exists $\lambda \in (-M, M)$ such that $u = u_\lambda$. Furthermore, observe that $T(M) > 0 > T(-M)$, which proves that a solution always exists. Moreover, $w_\lambda := \frac{\partial u_\lambda}{\partial \lambda}$ satisfies

$$w''_\lambda(x) = \frac{\partial g}{\partial u}(x, u_\lambda(x))w_\lambda(x), \quad w_\lambda(0) = 1, w'_\lambda(0) = a_0.$$

Thus we deduce the following result, more general than Theorem 1.1.

Proposition 3.1. *Let Φ be defined as the unique solution of the problem*

$$\Phi''(x) - \frac{\partial g}{\partial u}(x, 0)\Phi(x) = 0, \quad \Phi'(0) = a_0 \Phi(0) = a_0.$$

If $\Phi'(1) < a_1 \Phi(1)$, then (1.1)–(1.2) has at least three solutions for $\|A\|_\infty$ small.

Proof. In the previous setting observe that, when $A = 0$, $u_0 = 0$ and $e(0) = 1$; thus, $w_0 = \Phi$ and hence $T'(0) = \Phi'(1) - a_1 \Phi(1) < 0$. Since $T(-M) < 0 < T(M)$, it is deduced that T has three simple roots and the result follows by a continuity argument. \square

Proof of Theorem 1.1: Let $a := -\frac{\partial g}{\partial u}(x, 0)$, then using (1.5) we deduce, from Lemma 2.2, that $\text{sgn}(\Phi'(1) - a_1 \Phi(1)) = (-1)^k < 0$, because k is odd. Thus, the result follows from the previous proposition. \square

In the context of Proposition 3.1, let us now consider the opposite case $\Phi'(1) > a_1 \Phi(1)$, for which the existence of nontrivial roots of T for $A = 0$ cannot be ensured since $T'(0) > 0$. However, if for some λ it is verified that $\lambda T(\lambda) < 0$, then T has at least two zeros with the same sign of λ and, consequently, the problem has at least two nontrivial solutions. This fact shall be the main argument in our proof of Theorem 1.2, based on the existence of a positive and a negative λ as before. With this aim, let us firstly prove the following lemma.

Lemma 3.2. *Assume $\frac{\partial g}{\partial u}(x, 0) < 0$ and $a_1 \geq \frac{a_0}{a_0+1}$. Then (1.1)–(1.2) with $A = 0$ has at least a positive solution and a negative solution.*

Proof. Fix $\varepsilon > 0$ such that $\frac{\partial g}{\partial u}(x, u) \leq 0$ for $|u| \leq \varepsilon(a_0 + 1)$ and define $\alpha(x) := \varepsilon(a_0x + 1)$, then

$$\alpha''(x) \geq g(x, \alpha(x))$$

and

$$\alpha'(0) \geq a_0\alpha(0), \quad \alpha'(1) \leq a_1\alpha(1).$$

On the other hand, we may take for example $\beta(x) = e^{mx^2+c}$ with $m > 2a_1$ and $c \gg 0$, then

$$\beta''(x) \leq g(x, \beta(x))$$

and

$$\beta'(0) \leq a_0\beta(0), \quad \beta'(1) \geq a_1\beta(1).$$

Thus, the result is deduced from a straightforward adaptation of the method of upper and lower solutions (see e.g. [3]). The existence of a negative solution follows in a similar way. \square

Again, Theorem 1.2 shall be deduced from a more general result, namely the following proposition.

Proposition 3.3. *Let Φ be defined as before and assume that $\frac{\partial g}{\partial u}(x, 0) < 0$ and $a_1 > \frac{a_0}{a_0+1}$. If $\Phi'(1) > a_1\Phi(1)$, then (1.1)–(1.2) has at least five solutions for $\|A\|_\infty$ small.*

Proof. Fix $\tilde{a}_1 \in \left(\frac{a_0}{a_0+1}, a_1\right)$. From the previous lemma, there exist $u > 0 > v$ solutions of (1.1) with

$$u'(0) - a_0u(0) = v'(0) - a_0v(0) = 0, \quad u'(1) - \tilde{a}_1u(1) = v'(1) - \tilde{a}_1v(1) = 0.$$

It follows that $u = u_{\lambda_+}$ and $v = v_{\lambda_-}$ with $\lambda_+ = u(0) > 0$, $\lambda_- = v(0) < 0$. Since $u(1) > 0 > v(1)$ we deduce that

$$u'(1) - a_1u(1) = (\tilde{a}_1 - a_1)u(1) < 0 < (\tilde{a}_1 - a_1)v(1) = v'(1) - a_1v(1).$$

In other words,

$$T(\lambda_-) > 0 > T(\lambda_+)$$

and the result follows. \square

Proof of Theorem 1.2: Since $\lambda_2 > 0$, condition (1.5) with $k > 0$ implies that $\frac{\partial g}{\partial u}(\cdot, 0) < 0$. Moreover, since k is even we deduce from Lemma 2.2 that $\text{sgn}(\Phi'(1) - a_1\Phi(1)) = 1$ and the previous proposition applies. \square

4 Variational formulation

In this section, we introduce a variational formulation for problem (1.1)–(1.2), that allows to study multiplicity of solutions from a different point of view. To this end, let us define the functional $\mathcal{J} : H^1(0, 1) \rightarrow \mathbb{R}$ by

$$\mathcal{J}(u) := \int_0^1 \left(\frac{1}{2}u'(x)^2 + G(x, u(x)) + A(x)u(x) \right) dx + \frac{a_0}{2}u(0)^2 - \frac{a_1}{2}u(1)^2,$$

where $G(x, u) := \int_0^u g(x, s) ds$. It is readily seen that $\mathcal{J} \in C^1(H^1(0, 1), \mathbb{R})$, with

$$\begin{aligned} D\mathcal{J}(u)(v) &= \int_0^1 [u'(x)v'(x) + g(x, u(x))v(x) + A(x)v(x)] dx \\ &\quad + a_0u(0)v(0) - a_1u(1)v(1), \end{aligned}$$

and that $u \in H^1(0, 1)$ is a critical point of \mathcal{J} if and only if u is a classical solution of (1.1)–(1.2). From standard results (see e.g. [5]), \mathcal{J} is weakly lower semi-continuous. Moreover, write as before $a_1u(1)^2 - a_0u(0)^2 \leq C\|u\|_{L^2}^2 + \frac{1}{2}\|u'\|_{L^2}^2$ to conclude, from the superlinearity, that

$$\mathcal{J}(u) \geq \frac{1}{2}\|u\|_{H^1}^2 - K. \quad (4.1)$$

for some constant $K > 0$. Thus, the functional is coercive and hence achieves a global minimum. This proves, again, that the problem has at least one solution.

Remark 4.1. Observe that the existence of solutions holds, in fact, for arbitrary $A \in L^2(0, 1)$ and a weaker form of (1.3), namely: for every $M > 0$ there exists $K > 0$ such that

$$G(x, u) \geq Mu^2 - K \quad (4.2)$$

for all u .

In order to prove the existence of multiple solutions, let us firstly observe that \mathcal{J} satisfies the Palais–Smale condition, that is, if $\{\mathcal{J}(u_n)\}$ is bounded and $D\mathcal{J}(u_n) \rightarrow 0$ as $n \rightarrow \infty$, then $\{u_n\}$ has a convergent subsequence in $H^1(0, 1)$.

Indeed, let $\{u_n\} \subset H^1(0, 1)$ be a Palais–Smale sequence. Because \mathcal{J} is coercive, we may assume that $\{u_n\}$ converges weakly in $H^1(0, 1)$ and uniformly to some u . Since $\mathcal{J} \in C^1$ and $D\mathcal{J}(u_n)(u) \rightarrow 0$, we deduce

$$0 = D\mathcal{J}(u)(u) = \int_0^1 [u'(x)^2 + g(x, u(x))u(x) + A(x)u(x)] dx + a_0u(0)^2 - a_1u(1)^2.$$

Moreover, using the fact that $D\mathcal{J}(u_n)(u_n) \rightarrow 0$, it is seen that $\|u'_n\|_{L^2}^2 \rightarrow \|u'\|_{L^2}^2$. Since

$$\|u'_n - u'\|_{L^2}^2 = \|u'_n\|_{L^2}^2 + \|u'\|_{L^2}^2 - 2 \int_0^1 u'_n(x)u'(x) dx \rightarrow 0,$$

we conclude that $u_n \rightarrow u$ for the H^1 -norm.

Before stating our multiplicity result, for convenience we define, for each $K > 0$,

$$C_K := \max_{0 \leq x \leq 1, |u| \leq K} \frac{\partial g}{\partial u}(x, u).$$

In particular, condition (1.6) implies that $C_K < -\lambda_k$. Moreover, set $C_0 > 0$ as the best constant such that

$$\int_0^1 \left(\frac{1}{2}u'(x)^2 + A(x)u(x) \right) dx + \frac{a_0}{2}u(0)^2 \geq -C_0\|A\|_{L^2}^2$$

for all A and all u such that $u(1) = 0$.

Remark 4.2. The value of C_0 can be computed in the following way. Note that, for fixed A , the minimum of $\mathcal{F}(u) := \int_0^1 (\frac{1}{2}u'(x)^2 + A(x)u(x)) dx + \frac{a_0}{2}u(0)^2$ subject to the constraint $u(1) = 0$ is attained at

$$u_A(x) := \lambda(A)(x-1) + \int_x^1 \mathcal{A}(s) ds,$$

where $\mathcal{A}(x) := \int_x^1 A(s) ds$ and the Lagrange multiplier $\lambda(A)$ is given by

$$\lambda(A) := \frac{a_0 \int_0^1 \mathcal{A}(x) dx - \mathcal{A}(0)}{1 - a_0}.$$

Thus, C_0 is obtained by minimizing the functional $\mathcal{G}(A) := \mathcal{F}(u_A)$ under the constraint $\|A\|_{L^2} = 1$.

Thus, we deduce the following quantitative version of Theorem 1.3.

Theorem 4.3. Assume there exist $\eta > 0$ and $k \in \mathbb{N}$ such that, for $K = \eta \|\varphi_k\|_\infty$,

$$\frac{\lambda_k + C_K}{2} \eta^2 + \eta \|A\|_{L^2} + C_0 \|A\|_{L^2}^2 < \min G. \quad (4.3)$$

Then problem (1.1)–(1.2) has at least three solutions.

Proof. Define $X_1 := \text{span}\{\varphi_k\}$ and $X_2 := \{u \in H^1(0,1) : u(1) = 0\}$, then $H^1(0,1) = X_1 \oplus X_2$. On the one hand, from the previous definition we know that

$$\inf_{u \in X_2} \mathcal{J}(u) \geq \min G - C_0 \|A\|_{L^2}^2.$$

On the other hand, recall that the eigenfunction φ_k does not vanish at $x = 1$ and was chosen in such a way that $\varphi_k(0) > 0$ and $\|\varphi_k\|_{L^2} = 1$. Writing $G(x, u) = \frac{1}{2} \frac{\partial g}{\partial u}(x, \xi) u^2$, we may compute:

$$\begin{aligned} \mathcal{J}(\pm \eta \varphi_k) &= \int_0^1 \left(-\frac{\eta^2}{2} \varphi_k''(x) \varphi_k(x) + G(x, \pm \eta \varphi_k(x)) \pm \eta \varphi_k(x) A(x) \right) dx \\ &\leq \eta^2 \frac{\lambda_k + C_K}{2} + \eta \|A\|_{L^2} < \inf_{u \in X_2} \mathcal{J}(u) := \rho. \end{aligned}$$

From a well known linking theorem by Rabinowitz (see [6]), there exists a critical point u_1 such that $\mathcal{J}(u_1) \geq \rho > \min_{u \in H^1(0,1)} \mathcal{J}(u)$. This implies, in particular, that if u_0 is a global minimizer of \mathcal{J} then $u_0 \neq u_1$ and $u_0(1) \neq 0$. Let $s := \text{sgn}(u_0(1))$ and observe that there exists u_2 such that

$$\mathcal{J}(u_2) = \min_{\{u : su(1) \leq 0\}} \mathcal{J}(u).$$

It is clear that $\mathcal{J}(u_2) \leq \mathcal{J}((-1)^k s \eta \varphi_k) < \rho \leq \mathcal{J}(u_1)$. Again, this implies that $u_2 \neq u_1$ and that $u_2 \notin X_2$. It follows that u_2 is a local minimum of \mathcal{J} and $\text{sgn}(u_2(1)) \neq \text{sgn}(u_0(1))$. Thus, $u_2 \neq u_0$ and the proof is complete. \square

Proof of Theorem 1.3: Let $\eta := \frac{K}{\|\varphi_k\|_\infty}$, then condition (1.6) reads

$$\theta := \min G - \frac{C_K + \lambda_k}{2} \eta^2 > 0.$$

Thus, the result follows from Theorem 4.3, taking $A_1 := \frac{-\eta + \sqrt{\eta^2 + 4\theta C_0}}{C_0}$. \square

Proof of Theorem 1.4: From the computations in section 2, it is readily verified, on the one hand, that if we let $a_1 \rightarrow +\infty$ then $\lambda_1 = -r^2$ with $\frac{r}{a_1} \rightarrow 1^+$. In particular, when $a_1 \gg 0$,

$$\varphi_1(x) = a \left(e^{rx} + \frac{r - a_0}{r + a_0} e^{-rx} \right)$$

with $a \simeq \sqrt{\frac{2r}{e^{2r}-1}}$ and $r \simeq a_1 \gg 0$. This implies

$$\|\varphi_1\|_\infty = \varphi_1(1) \simeq ae^r \simeq \sqrt{2a_1}$$

for $a_1 \gg 0$. In particular, fixing an arbitrary $K > 0$ it follows that condition (1.6) with $k = 1$ is satisfied for a_1 sufficiently large. Furthermore, setting η and θ as in the previous proof it is verified that $\theta = O(a_1)$ and, consequently, $A_1 \rightarrow +\infty$ as $a_1 \rightarrow +\infty$. On the other hand, observe that if $a_1 < \frac{a_0}{a_0+1}$ then $\lambda_1 = r^2$, where r is the first positive solution of the equation

$$-r \sin r + a_0 \cos r = a_1 \left(\cos r + a_0 \frac{\sin r}{r} \right).$$

Thus, letting $a_1 \rightarrow 0$, it is seen that $\lambda_1 \rightarrow r_1^2$. Hence, if a_1 is sufficiently small then $\frac{\partial g}{\partial u}(x, u) > -\lambda_1$ for all x and all u . Uniqueness follows then from Theorem 2.2 in [1]. \square

As a final remark, it is worth mentioning that Theorem 1.4 is not directly deduced from the above shooting arguments. On the one hand, when a_1 is large, it is readily verified that (1.5), as well as the weaker condition of Proposition 3.1, do not necessarily hold. Furthermore, even if one of these conditions holds, it is not clear how to get rid of the smallness condition on A . Finally, it is worth mentioning that the shooting operator T depends on a_1 , which makes it difficult to handle when a_1 gets large. On the other hand, when a_1 is small it is possible to prove the existence of multiple solutions for some specific choices of g . For example, it suffices to observe that λ_2 tends, as $a_1 \rightarrow 0$, to some $r_2^2 > r_1^2$. Thus, fixing g such that

$$-r_1^2 > \frac{\partial g}{\partial u}(\cdot, 0) > -r_2^2,$$

the existence of three solutions follows from Theorem 1.1 if a_1 and $\|A\|_\infty$ are small enough. Other possible multiplicity conditions, however, require that a_1 is not too small: for example, in Theorem 1.2, it is not clear whether or not the condition on a_1 can be relaxed.

Acknowledgements

This work was partially supported by project UBACyT 20020120100029BA.

References

- [1] P. AMSTER, M. P. KUNA, On exact multiplicity for a second order equation with radiation boundary conditions, *submitted*, 2016.
- [2] P. AMSTER, M. K. KWONG, C. ROGERS, A Painlevé II model in two-ion electrodiffusion with radiation boundary conditions, *Nonlinear Anal. Real World Appl.* **16**(2014), No. 16, 120–131. [MR3123805](#); [url](#)

- [3] C. DE COSTER, P. HABETS, Upper and lower solutions in the theory of ODE boundary value problems: classical and recent results, in: F. Zanolin (ed.), *Non-linear analysis and boundary value problems for ordinary differential equations (Udine)*, CISM Courses and Lectures, Vol. 371, Springer, Vienna, 1996, pp. 1–78. [MR1465239](#)
- [4] A. C. LAZER, Application of a lemma on bilinear forms to a problem in nonlinear oscillation, *Amer. Math. Soc.* **33**(1972), No. 33, 89–94. [MR0293179](#); [url](#)
- [5] J. MAWHIN, M. WILLEM, *Critical point theory and Hamiltonian systems*, Springer-Verlag, New York, 1989. [MR982267](#); [url](#)
- [6] P. RABINOWITZ, Some minimax theorems and applications to partial differential equations, in: *Nonlinear analysis (collection of papers in honor of Erich H. Rothe)*, Academic Press, New York, 1978, pp. 161–177. [MR0501092](#)