

УДК 512.555+517.982

BANACH LATTICES OF CONTINUOUS SECTIONS¹

A. G. Kusraev, S. N. Tabuev

The aim of this note is to outline some application of ample continuous Banach bundles to the theory of Banach lattices.

Mathematics Subject Classification (2000): 06F25, 46A40.

Key words: Banach lattice, continuous Banach bundle, section, injective Banach lattice.

1. Introduction

The study of Banach lattices in terms of sections of continuous Banach bundles has been started by Giertz [1, 2]. Later Gutman create the theory of ample (or complete) continuous Banach bundles [3] and measurable Banach bundles admitting lifting [4]. A portion of the Gutman's theory was specified in the case of bundles of measurable Banach lattices by Ganiev [5] and Kusraev [6]. The aim of this short note is to outline some additional possibilities of applying ample Banach bundles to the theory of Banach lattices. Recall some definitions.

A *bundle of Banach lattices* over a set Q is a mapping \mathcal{X} defined on Q and sending every point $q \in Q$ to a Banach lattice $\mathcal{X}(q) := (\mathcal{X}(q), \|\cdot\|_q)$. Each space $\mathcal{X}(q)$ of a bundle \mathcal{X} is called its *stalk* over q . A mapping u defined on a nonempty subset $D \subset Q$ is called a *section* over D , if $u(q) \in \mathcal{X}(q)$ for every $q \in D$. A section over Q is called *global*. If Q is endowed with some topology we call sections over comeager subsets of Q *almost global*.

Let $S(Q, \mathcal{X})$ stands for the set of all global sections of \mathcal{X} , endowed with the structure of a vector lattice by letting $u \leq v \iff u(q) \leq v(q) (\forall q \in Q)$, and $(\alpha u + \beta v)(q) = \alpha u(q) + \beta v(q)$ ($q \in Q$), where $\alpha, \beta \in \mathbb{R}$ and $u, v \in S(Q, \mathcal{X})$. For each section $u \in S(Q, \mathcal{X})$ we define its point-wise norm by $\|u\| : q \mapsto \|u(q)\|_{\mathcal{X}(q)}$ ($q \in Q$). A set of sections $\mathcal{U} \subset S(Q, \mathcal{X})$ is called stalk-wise dense in \mathcal{X} if the set $\{u(q) : u \in \mathcal{U}\}$ is dense in $\mathcal{X}(q)$ for every $q \in Q$.

2. Continuous bundles of Banach lattices

Let Q be a topological space and \mathcal{X} be a bundle of Banach lattices over Q . A set of global sections $\mathcal{C} \subset S(Q, \mathcal{X})$ is called a *continuity structure* on \mathcal{X} , if it satisfy the conditions:

- \mathcal{C} is a vector lattice, i. e. $\alpha c_1 + \beta c_2 \in \mathcal{C}$, $|c| \in \mathcal{C}$ for all $\alpha, \beta \in \mathbb{R}$ and $c_1, c_2 \in \mathcal{C}$;
- the point-wise norm $\|c\| : Q \rightarrow \mathbb{R}$ is continuous for every $c \in \mathcal{C}$;
- \mathcal{C} is stalk-wise dense in \mathcal{X} .

If \mathcal{C} is a continuity structure on \mathcal{X} then the pair $(\mathcal{X}, \mathcal{C})$ is called a *continuous bundle of Banach lattices over Q* . More details see in [3] and [7]. Below $(\mathcal{X}, \mathcal{C})$ stands for a continuous bundle of Banach lattices over Q . We say that a section $u \in S(D, \mathcal{X})$ over $D \subset Q$ is \mathcal{C} -continuous at the point $q \in D$ if the function $\|u - c\|$ is continuous at q for every $c \in \mathcal{C}$. A section $u \in S(D, \mathcal{X})$ is \mathcal{C} -continuous if it is \mathcal{C} -continuous at every $q \in D$.

© 2012 Kusraev A. G., Tabuev S. N.

¹The study was supported by The Ministry of education and science of Russian Federation, project 8210; by a grant from the Russian Foundation for Basic Research, project 12-01-00623-a.

Lemma. *Let $(\mathcal{X}, \mathcal{C})$ be a continuous bundle of Banach lattices over Q . The set of all \mathcal{C} -continuous sections over $D \subset Q$ is a vector lattice.*

◁ It is obvious that the set of all \mathcal{C} -continuous sections is a vector space. Ensure that if a section u is \mathcal{C} -continuous then so is $|u| : q \mapsto |u(q)|$ ($q \in D$). It is sufficient to prove that the function $\| |u| - c \| : Q \rightarrow \mathbb{R}$ is continuous at an arbitrary $q \in D$, for every $c \in \mathcal{C}$. Put $\lambda := \| |u|(q) - c(q) \|$. We have to prove that, given $q \in D$ and $\epsilon > 0$, one can choose a neighborhood U of q such that $\lambda - \epsilon < \| |u|(p) - c(p) \| < \lambda + \epsilon$ for every $p \in U$.

Select a section $v \in \mathcal{C}$ satisfying $\|u(q) - v(q)\| < \epsilon/2$. Observe that $\| |u|(q) - |v|(q) \| \leq \|u(q) - v(q)\| < \epsilon/2$. Taking into consideration the continuity of the function $\| |v| - c \|$ and the estimate $\| |v|(q) - c(q) \| \leq \| |u|(q) - c(q) \| + \| |u|(q) - |v|(q) \| < \lambda + \epsilon/2$ we can find a neighborhood U_1 of q with $\| |v|(p) - c(p) \| < \epsilon/2$ for all $p \in U_1$.

Similarly, the estimate $\| |v|(q) - c(q) \| \geq \| |u|(q) - c(q) \| - \| |u|(q) - |v|(q) \| \geq \lambda - \epsilon/2$ implies that $\| |v|(p) - c(p) \| \geq \lambda - \epsilon/2$ ($p \in U_2$) for some neighborhood U_2 of q . Now, for all $p \in U := U_1 \cap U_2$ we can easily deduce

$$\begin{aligned} \lambda - \epsilon &= (\lambda - \epsilon/2) - \epsilon/2 < \| |v|(p) - c(p) \| - \| |u|(p) - |v|(p) \| \leq \| |u|(p) - c(p) \|, \\ \| |u|(p) - c(p) \| &\leq \| |v|(p) - c(p) \| + \| |u|(p) - |v|(p) \| < (\lambda + \epsilon/2) + \epsilon/2 = \lambda + \epsilon. \triangleright \end{aligned}$$

3. Banach lattices of sections

Suppose that Q is a nonempty Stonean space (\equiv extremally disconnected and compact Hausdorff space). Consider a continuous Banach bundle \mathcal{X} over Q . If u is an almost global section of the bundle \mathcal{X} then the function $q \mapsto \|u(q)\|_q$ is defined and continuous on a comeager set $\text{dom}(u) \subset Q$. Consequently, there exists a unique function $|u| \in C_\infty(Q)$ such that $|u|(q) = \|u(q)\|_q$ ($q \in \text{dom}(u)$).

In the set of almost global sections $\mathfrak{M}(Q, \mathcal{X})$ we can define an equivalence relation by letting $u \sim v$ if $u(q) = v(q)$ whenever $q \in \text{dom}(u) \cap \text{dom}(v)$. Then equivalent u and v we have $|u| = |v|$; therefore, we may define $|\tilde{u}| := |u|$, where \tilde{u} is the coset of the almost global section u . Denote by $C_\infty(Q, \mathcal{X})$ the quotient space $\mathfrak{M}(Q, \mathcal{X}) / \sim$.

In each coset \tilde{u} , there exists a unique section $\bar{u} \in \tilde{u}$ such that $\text{dom}(v) \subset \text{dom}(\bar{u})$ for all $v \in \tilde{u}$. The section \bar{u} is called extended. The space $C_\infty(Q, \mathcal{X})$ can be represented also as the space of all extended almost global sections of the bundle \mathcal{X} , see [3]. The set $C_\infty(Q, \mathcal{X})$ can be naturally equipped with the structure of lattice-normed lattice. For instance, the element $\tilde{u} + \tilde{v}$ is defined as the coset of the almost global section $q \mapsto u(q) + v(q)$ ($q \in \text{dom}(u) \cap \text{dom}(v)$). If E is an order ideal in $C_\infty(Q)$ then we assign $E(\mathcal{X}) := \{u \in C_\infty(Q, \mathcal{X}) : |u| \in E\}$.

Recall that a *Banach–Kantorovich space* over a Dedekind complete vector lattice E is a vector space X with a decomposable norm $|\cdot| : X \rightarrow E_+$ which is norm complete with respect to order convergence in E . Decomposability means that, given $e_1, e_2 \in E_+$ and $x \in X$ with $|x| = e_1 + e_2$, there exist $x_1, x_2 \in X$ such that $x = x_1 + x_2$ and $|x_k| = e_k$ ($k := 1, 2$). If a Banach–Kantorovich space is in addition a vector lattice with monotone norm then it's called a *Banach–Kantorovich lattice*. A Banach–Kantorovich lattice X can be endowed with a scalar norm $x \mapsto \| |x| \| := \| |\cdot| \|_E$, whenever E is a Banach lattice. The following result see in [3, 7].

Theorem 1. *Let \mathcal{X} be a continuous bundle of Banach lattices over a Stonean space Q . Then $C_\infty(Q, \mathcal{X})$ is a Banach–Kantorovich lattice over $C_\infty(Q)$. If E is an order ideal in $C_\infty(Q)$ then $(E(\mathcal{X}), |\cdot|)$ is a Banach–Kantorovich lattice over E . If, in addition, E is Banach lattice, then $(E(\mathcal{X}), \| |\cdot| \|)$ is a Banach lattice.*

◁ We need only to put together the ‘Banach part’, given in [3] and [7, Theorem 2.4.7], and the above Lemma. ▷

Theorem 2. *Every Banach–Kantorovich lattice X over an order dense ideal $E \subset C_\infty(Q)$ is isometrically lattice isomorphic to $E(\mathcal{X})$ for some complete continuous bundle \mathcal{X} of Banach lattices over Q . Moreover, such a bundle \mathcal{X} is unique to within isometrically lattice isomorphism.*

◁ The ‘Banach part’ follows again from [3] (see also [7, 2.4.10]). The rest is easily deduced on using the above Lemma. ▷

Let Q be the Stone space of the Boolean algebra $B(\Omega)$ and $\tau : \Omega \rightarrow Q$ is the canonical immersion of Ω into Q corresponding to a fixed lifting τ of $L^\infty(\Omega)$. Let \mathcal{Y} be a complete continuous bundle of Banach lattices over Q and $\mathcal{X} = \mathcal{Y} \circ \tau$. If \mathcal{C} is a continuous structure in \mathcal{Y} , then the set $\mathcal{C} \circ \tau$ is a measurability structure in \mathcal{X} . The composite $v \circ \tau$ is a measurable section of \mathcal{X} for every $v \in C_\infty(Q, \mathcal{Y})$, see [2, 1.2.7, 1.4.9, 2.5.8]. Let $C(Q, \mathcal{X})$ stands for the set of all global continuous sections of \mathcal{X} . The following result may be considered as a bridge between continuous and measurable bundles of Banach lattices.

Theorem 3. *Let (Ω, Σ, μ) be a measurable space with the direct sum property. The mapping $v \mapsto (v \circ \tau)^\sim$ is isometric lattice isomorphism of Banach–Kantorovich lattices $C_\infty(Q, \mathcal{Y})$ and $L^0(\Omega, \mathcal{X})$, associated with the isomorphism $(e \mapsto (e \circ \tau)^\sim) : C_\infty(Q) \rightarrow L^0(\Omega)$. The image of $C(Q, \mathcal{Y})$ under this isomorphism is $L^\infty(\Omega, \mathcal{X})$.*

◁ The ‘Banach part’ can be found in [4] (see also [2, 2.5.9]). The remaining is obvious. ▷

REMARK 1. The theory of ample continuous bundles of Banach lattices is parallel to that of liftable Banach bundles presented in [6]. In particular, the results from [6, Theorems 2.9, 2.10, 3.3] have their counterparts for ample continuous bundles of Banach lattices.

4. Representation of injective Banach lattices

A real Banach lattice X is said to be *injective* if, for every Banach lattice Y , every closed vector sublattice $Y_0 \subset Y$, and every positive linear operator $T_0 : Y_0 \rightarrow X$ there exists a positive linear extension $T : Y \rightarrow X$ with $\|T_0\| = \|T\|$. This concept was introduced by Lotz [8]. Important contributions are due to Cartwright [9] and Haydon [10]. A new source of insight into the structure of injectives is a Boolean-valued approach, see [11, 12].

A band projection π in a Banach lattice X is said to be an *M-projection* if $\|x\| = \max\{\|\pi x\|, \|\pi^\perp x\|\}$ for all $x \in X$, where $\pi^\perp := I_X - \pi$. The collection of all *M-projections* forms a subalgebra $\mathbb{M}(X)$ of the Boolean algebra of all band projections $\mathbb{P}(X)$ in X . The closed *f*-subalgebra in the center $\mathcal{Z}(X)$ generated by $\mathbb{M}(X)$ is denoted by $\mathcal{Z}_m(X)$.

A positive operator $T : X \rightarrow F$ is said to have the *Levi property* if $T(X)^{\perp\perp} = F$ and $\sup x_\alpha$ exists in X for every increasing net $(x_\alpha) \subset X_+$, provided that the net (Tx_α) is order bounded in F . A *Maharam operator* is an order continuous order intervals preserving ($\equiv T([0, x]) = [0, Tx]$ for all $x \in X_+$) operator. An operator $T : X \rightarrow Y$ is called *lattice \mathbb{B} -isometry*, if it is a lattice isometry and $b \circ T = T \circ b$ for all $b \in \mathbb{B}$.

Theorem 4. *If Φ is a strictly positive Maharam operator with the Levi property taking values in a Dedekind complete AM-space Λ with unit and $\|x\| = \|\Phi(|x|)\|_\infty$ ($x \in L^1(\Phi)$), then $(L^1(\Phi), \|\cdot\|)$ is an injective Banach lattice with $\mathbb{M}(L^1(\Phi)) = \mathbb{P}(\Lambda)$. Conversely, any injective Banach lattice X is lattice \mathbb{B} -isometric to $(L^1(\Phi), \|\cdot\|)$ for some strictly positive Maharam operator Φ with the Levi property taking values in a Dedekind complete AM-space Λ with unit, where $\mathbb{B} = \mathbb{M}(L^1(\Phi)) = \mathbb{P}(\Lambda)$.*

◁ See [12]; details can be found in [11]. ▷

Theorem 5. *Every injective Banach lattice X with $\Lambda = \mathcal{L}_m(X) = C(Q)$ and $\mathbb{B} := \mathbb{P}(\Lambda)$ is lattice \mathbb{B} -isometric to $\Lambda(\mathcal{X})$ for some complete continuous bundle \mathcal{X} of Banach lattices over Q such that all stalks $\mathcal{X}(q)$ ($q \in Q$) are AL-spaces. Moreover, such a bundle \mathcal{X} is unique to within isometrically lattice isomorphism.*

◁ The proof consists of a combination of the representation Theorems 2 and 4. ▷

REMARK 2. This result was proved essentially by Gierz [1, 2] and Haydon [10]. The above approach enables us to settle also the uniqueness problem.

References

1. Gierz G. Darstellung von Banachverbänden durch Schnitte in Bündeln // Mitt. Math. Sem. Univ. Giessen.—1977.—Vol. 125.
2. Gierz G. Bundles of Topological Vector Spaces and Their Duality.—Berlin etc.: Springer-Verlag, 1982.
3. Gutman A. E. Banach bundles in the theory of lattice-normed spaces. I. Continuous Banach bundles // Siberian Adv. Math.—1993.—Vol. 3, № 3.—P. 1–55.
4. Gutman A. E. Banach bundles in the theory of lattice-normed spaces. II. Measurable Banach bundles // Siberian Adv. Math.—1993.—Vol. 3, № 4.—P. 8–40.
5. Ganiev I. G. Measurable bundles of lattices and their applications // Studies on functional analysis and its applications.—Moscow: Nauka, 2006.—P. 9–49.
6. Kusraev A. G. Measurable bundles of Banach lattices // Positivity.—2010.—Vol. 14.—P. 785–799.
7. Kusraev A. G. Dominated Operators.—Dordrecht: Kluwer, 2000.—446 p.
8. Lotz H. P. Extensions and liftings of positive linear mappings on Banach lattices // Trans. Amer. Math. Soc.—1975.—Vol. 211.—P. 85–100.
9. Cartwright D. I. Extension of positive operators between Banach lattices // Memoirs Amer. Math. Soc.—1975.—Vol. 164.—P. 1–48.
10. Haydon R. Injective Banach lattices // Math. Z.—1977.—Vol. 156.—P. 19–47.
11. Kusraev A. G. Boolean Valued Analysis Approach to Injective Banach Lattices.—Vladikavkaz: Southern Math. Inst. VSC RAS, 2011.—28 p.—(Preprint № 1).
12. Kusraev A. G. Boolean-valued analysis and injective Banach lattices // Doklady Ross. Akad. Nauk.—2012.—Vol. 444, № 2.—P. 143–145; Engl. transl.: Doklady Mathematics.—2012.—Vol. 85, № 3.—P. 341–343.

Received Desember 9, 2012.

ANATOLY G. KUSRAEV
Southern Mathematical Institute
Vladikavkaz Science Center of the RAS, Director
Russia, 362027, Vladikavkaz, Markus street, 22
E-mail: kusraev@smath.ru

SOSLAN N. TABUEV
Southern Mathematical Institute
Vladikavkaz Science Center of the RAS, Researcher
Russia, 362027, Vladikavkaz, Markus street, 22
E-mail: soslan@tabuev.com

БАНАХОВЫ РЕШЕТКИ НЕПРЕРЫВНЫХ СЕЧЕНИЙ

Кусраев А. Г., Табуев С. Н.

Заметка представляет собой набросок некоторых приложений пространственных банаховых расслоений к теории банаховых решеток.

Ключевые слова: банахова решетка, непрерывное банахово расслоение, сечение, инъективная банахова решетка.